

Near-Infrared Identification of the Dimmest X-ray Sources on the Galactic Plane with the ESO/NTT SOFI Camera

KEN EBISAWA^{1,2}, V. BECKMANN¹, T. J.-L. COURVOISIER^{1,3}, P. DUBATH¹, H. KANEDA⁴, Y. MAEDA⁴, S. YAMAUCHI⁵ AND E. NISHIHARA⁶

¹ INTEGRAL Science Data Centre, Chemin d'Écogia 16, CH-1290 Versoix, Switzerland

² Laboratory for High Energy Astrophysics, NASA/GSFC, Greenbelt, MD 20771, USA

³ Geneva Observatory, Chemin des Maillettes 51, CH-1290 Sauverny, Switzerland

⁴ Institute of Space and Astronautical Science, Yoshinodai 3-1-1, Sagami-hara, Kanagawa 229-8510, Japan

⁵ Faculty of Humanities and Social Sciences, Iwate University, Ueda 3-18-34, Morioka, Iwate 020-8550, Japan

⁶ Gunma Astronomical Observatory, 6860-86 Nakayama, Takayama-mura, Agatsuma-gun, Gunma 377-0702, Japan

1. Introduction

We have carried out a deep X-ray observation on a typical Galactic plane region with the Chandra ACIS-I instrument with unprecedented sensitivity and spatial resolution (Ebisawa et al. 2001), and detected 274 unidentified X-ray point sources in the ~ 500 arcmin² region. In order to identify these new X-ray sources, we have carried out a near infrared follow-up observation using ESO/NTT (Tarengi and Wilson 1989) SOFI infrared camera on 2002/7/28 and 2002/7/29.

We have made a mosaic observation to cover the large Chandra field of view ($17' \times 17'$) with SOFI ($4.94' \times 4.94'$). We have chosen 7 “A” fields and surrounding 9 “B” fields to cover the central part of the Chandra field (Figure 1). Exposure time for J , H and K_s bands for an A-field is 10, 10 and 14 minutes, and those for a B-field is 5, 5 and 7.47 minutes respectively. The seeing was best in the first night ($\sim 0.6''$), whereas in the second night it was moderate ($\sim 1.5''$) when we observed the remaining fields.

2. Preliminary Results

Quick look analysis suggests we have reached the limiting magnitude ~ 20 mag for all the three bands in both A and B fields. In Figure 2, we show a zoomed K_s -band image of a part of field A2. The Chandra sources detected below ~ 2 keV are marked with red, and those detected above ~ 3 keV are marked with blue. All the five soft X-ray sources in this field have near-infrared counterparts, while the counterpart of the hard X-ray source (#151) is not found.

Most of the soft X-ray sources are considered to be nearby active stars, while hard X-ray sources are primarily background AGNs. From the J , H and K_s band color analysis, we plan to investigate the origin of these dim X-ray sources.




Fig. 1. SOFI fields of view on the Chandra image. Red ones are priority “A”, while green ones are priority “B”.

References

- Ebisawa, K. et al. ; 2001, Science 293, 1633
Tarengi, M. and Wilson, R. N.; 1989, SPIE, 1114, 302




Fig. 2. A SOFI K_s -band image of a part of field A2. Red circles indicate positions of the detected soft X-ray sources, while the blue one is a hard X-ray source.